Author Index

Alinovi, R., see Ghiggeri, G. M. 347
Antia, F. D., see Maa, Y.-F. 331
Armstrong, D. W.
——, Faulkner Jr., J. R. and Han, S. M.
Use of hydroxypropyl- and hydroxyethyl-derivatized β-cyclodextrins for the thin-layer chromatographic separation of enantiomers and diastereomers 323
Barkley, R. M., see McNamara, E. A. 75
Base, J., see Hanis, T. 443
Beckers, J. L.
——, Verheggen, Th. P. E. M. and Everaerts, F. M.
Use of a double-detector system for the measurement of mobilities in zone electrophoresis 591
——, see Verheggen, Th. P. E. M. 615
Beerahee, M., see Jack, D. B. 257
Berg, J. H. M. van den, see Tipker, J. 227
Bergamaschi, E., see Ghiggeri, G. M. 347
Bergh-Swart, J. K. van den, see Tipker, J. 227
Boček, P., see Foret, F. 601
Borówko, M.
—— and Jaroniec, M.
Studies of adsorption and partition effects in liquid chromatography with mixed mobile phases 131
Brinkman, U. A. Th., see Farjam, A. 419
Brooksbank, R. A., see Jackson, P. J. 359
Brown, J., see O'Flynn, M. A. 469
Brown, P. R., see Fadool, D. A. 491
Bruner, F., see Mangani, F. 527
Burford, R. P., see Haken, J. K. 37
Burke, J. T., see Krstulović, A. M. 477
Candiano, G., see Ghiggeri, G. M. 347
Causon, R. C., see O'Flynn, M. A. 469
Cobb, S. J., see Fadool, D. A. 491
Cooper, E. H., see Jackson, P. J. 359
Crescentini, G., see Mangani, F. 527
Croteau, R. B., see Satterwhite, D. M. 61
Crumley, F. G., see Xu, S. 377
Davankov, V. A.
——, Kurganov, A. A. and Ponomareva, T. M.
Enantioselectivity of complex formation in ligand-exchange chromatographic systems with chiral stationary and/or chiral mobile phases 309
David, P. A.
—— and Novotny, M.
Separation of α-keto acids by capillary supercritical fluid chromatography as their quinoxalinol derivatives 623
Dawkins, J. V.
——, Gabbott, N. P., Lloyd, L. L., McConville, J. A. and Warner, F. P.
Reversed-phase high-performance liquid chromatography with a C_{18} polyacrylamide-based packing 145
De Jong, G. J., see Farjam, A. 419
De Leenheer, A. P., see Nelis, H. J. C. F. 535
Deml, M., see Foret, F. 601
Deyl, Z., see Hanis, T. 443
Durand, A., see Krstulović, A. M. 477
Egestad, B., see Marshall, H.-U. 459
El Rassi, Z., see Maa, Y.-F. 331
Everaerts, F. M., see Beckers, J. L. 591
——, see Verheggen, Th. P. E. M. 615
Fadool, D. A.
——, Cobb, S. J., Kass-Simon, G. and Brown, P. R.
Liquid chromatographic procedures for the analysis of compounds in the serotenergic and octoptamine pathways of lobster hemolymph 491
Farjam, A.
——, De Jong, G. J., Frei, R. W., Brinkman, U. A. Th., Haasnoot, W., Hamers, A. R. M., Schilt, R. and Huf, F. A.
Immunoaffinity pre-column for selective on-line sample pre-treatment in high-performance liquid chromatography determination of 19-nortestosterone 419
Faulkner Jr., J. R., see Armstrong, D. W. 323
Fisher, D. H.
—— and Giese, R. W.
Determination of 5-methylcytosine in DNA by gas chromatography–electron-capture detection 51
Floyd, T. R.
——, Sagliano Jr., N. and Hartwick, R. A.
Analysis of short alkylsilane bonded silica gel high-performance liquid chromatographic stationary phases using hydrofluoric acid digestion and headspace analysis by capillary gas chromatography 43
Foret, F.
——, Deml, M. and Boček, P.
Capillary zone electrophoresis. Quantitative study of the effects of some dispersive processes on the separation efficiency 601
Fouchet, M. H., see Krstulović, A. M. 477
Frei, R. W., see Farjam, A. 419
Gabbott, N. P., see Dawkins, J. V. 145

Ghiggeri, G. M.
——, Candiano, G., Ginevri, F., Mutti, A., Bergamaschi, E., Alinovi, R. and Righetti, P. G.
Hydrophobic interaction of Alcian Blue with soluble and erythrocyte membrane proteins 347
Ghodbane, S.
—— and Guiochon, G.
Effect of mobile phase flow-rate on the recoveries and production rates in overloaded elution chromatography. A theoretical study 209
Giese, R. W., see Fisher, D. H. 51
——, see Trainor, T. M. 369
Gillet, G., see Krstulović, A. M. 477
Ginevri, F., see Ghiggeri, G. M. 347
Glikberg, S., see Shohat, S. 503
Görög, S.
——, Herényi, B., Nyéki, O., Schön, I. and Kisfaludy, L.
Determination of impurity profiles in drugs and related materials. II. Detection and quantification of a diastereomeric impurity in the peptide RGH-0205 (Arg-Lys-Asp) 317
Golombek, F.
—— and Schwedt, G.
Methyl green-coated column for separation of inorganic anions by ion chromatography 283
Goto, J.
——, Miura, H., Inada, M., Nambara, T., Nagakura, R. and Suzuki, H.
Studies on steroids. CCXXXVIII. Determination of bile acids in liver tissue by gas chromatography–mass spectrometry with negative ion chemical ionization detection 119
Green, G., see Marshall, H.-U. 459
Groen, C. P., see Tipker, J. 227
Grushka, E., see Shohat, S. 503
Guiochon, G., see Ghodbane, S. 209
Haasnoot, W., see Farjam, A. 419
Hais, I. M.
Tswett's letter to Claparède on tropism and taxes 5
Haken, J. K.
—— and Smith, R. J.
Dispersion and selectivity indices of alkyl and alkenyl benzenes 31
——, Harahap, N. and Burford, R. P.
Quantitative determination of dicarboxylic acids and polyols in silicone–polyester resins 37
Hamers, A. R. M, see Farjam , A. 419
Han, S. M., see Armstrong, D. W. 323
Hanis, T.
——, Smrz, M., Klir, P., Macek, K., Klima, J., Base, J. and Deyl, Z.
Determination of fatty acids as phenacyl esters in rat adipose tissue and blood vessel walls by high-performance liquid chromatography 443
Harahap, N., see Haken, J. K. 37
Hartwick, R. A., see Floyd, T. R. 43
Hawker, J. L., see Jack, D. B. 257
Herényi, B., see Görög, S. 317
Hongxin, H., see Peichang, L. 175
Horvath, Cs., see Maa, Y.-F. 331
Huang, T.
——, Wall, F. and Kabra, P.
Improved solid-phase extraction and liquid chromatography with electrochemical detection of urinary catecholamines and 5-S-L-cysteinyl-L-dopa 409
Huf, F. A., see Farjam, A. 419
Inada, M., see Goto, J. 119
Issaq, H. J.
——, Williams, D. G., Schultz, N. and Saavedra, J. E.
High-performance liquid chromatography separations of nitrosamines. III. Conformers of N-nitrosamino acids 511
Jack, D. B.
——, Hawker, J. L., Rooney, L., Beerahee, M. Lobo, J. and Patel, P.
Measurement of the distribution coefficients of several classes of drug using reversed-phase thin-layer chromatography 257
Jackson, P. J.
——, Turner, R., Keen, J. N., Brooksbank, R. A. and Cooper, E. H.
Purification and partial amino acid sequence of human urine protein. I. Evidence for homology with rabbit uteroglobin 359
Janča, J.
—— and Nováková, N.
Retention in sedimentation–flotation focusing field-flow fractionation using a step density gradient 549
Jaroniec, M., see Borówko, M. 131
Jelínek, I., see Snopek, J. 571
Jellum, E.
Chromatography for diagnosis of metabolic diseases 435
Johansson, M.
—— and Westerlund, D.
Effects of system peaks in ion-pair reversed-phase liquid chromatography for noscapine and metabolites 241
Jong, G. J. de, see Farjam, A. 419
Kabra, P., see Huang, T. 409
Kageyama, S., see O'Flynn, M. A. 469

Kass-Simon, G., see Fadool, D. A. 491
Keen, J. N., see Jackson, P. J. 359
Kersten, B. R.
—— and Poole, C. F.
Considerations for using the solvent selectivity triangle approach for stationary phase characterization 191
Kisfaludy, L., see Görög, S. 317
Klima, J., see Hanis, T. 443
Klir, P., see Hanis, T. 443
Krstulović, A. M.
——, Fouchet, M. H., Burke, J. T., Gillet, G. and Durand, A.
Direct enantiomeric separation of betaxolol with applications to analysis fo bulk drug and biological samples 477
Krull, I. S., see Lookabaugh, M. 295
Kuksis, A., see Myher, J. J. 93
Kurganov, A. A., see Davankov, V. A. 309
Ladizinsky, G., see Milo, J. 563
Lederer, M.
Some considerations on the 'charge' on a metal ion in ion-exchange Chromatography 265
Leenheer, A. P., De, see Nelis, H. J. C. F. 535
Levy, A., see Milo, J. 563
——, see Weissenberg, M. 485
Levy, E. C., see Weissenberg, M. 485
Lin, J.-T.
—— and Stafford, A. E.
Reveresed-phase C_{18} and normal-phase silica high-performance liquid chromatography of gibberellins and their methyl esters 519
Lloyd, L. L., see Dawkins, J. V. 145
Lobo, J., see Jack, D. B. 257
Lookabaugh, M.
—— and Krull, I. S.
Determination of nitrite and nitrate by reversed-phase high-performance liquid chromatography using on-line post-column photolysis with ultra violet absorbance and electrochemical detection 295
Maa, Y.-F.
——, Antia, F. D., El Rassi, Z. and Horváth, Cs.
Mixed-bed ion-exchange columns for protein high-performance liquid chromatography 331
McConville, J. A., see Dawkins, J. V. 145
Macek, K., see Hanis, T. 443
McNamara, E. A.
——, Montzka, S. A., Barkley, R. M. and Sievers, R. E.
Superconductor metal oxide catalyst in a chemiluminescence chromatography detector 75
Mangani, F.
——, Crescentini, G., Palma, P. and Bruner, F.
Performance of graphitized carbon black cartridges in the extraction of some organic priority pollutants form water 527
Marai, L., see Myher, J. J. 93
Marshall, H.-U.
——, Green, G., Egestad, B. and Sjövall, J.
Isolation of bile acid glucosides and N-acetylglucosaminides from human urine by ion-exchange chromatography and reversed-phase high-performance liquid chromatography 459
Martin, M.
——, Thevenon, G. and Tchapla, A.
Comparison of retention mechanisms of homologous series and triglycerides in non-aqueous reversed-phase liquid chromatography 157
Martire, D. E.
Unified theory of adsorption chromatography: gas, liquid and supercritical fluid mobile phases 17
Matsuura, S., see Niwa, T. 85
Michalec, Č.
—— and Ranný, M.
Potential application of thin-layer chromatography and thin-layer chromatography with flame ionization detection of cholestanol in the diagnosis of cerebrotendinous xanthomatosis 543
Milo, J.
——, Levy, A., Palevitch, D. and Ladizinsky, G.
High-performance liquid chromatographic analysis of alkaloid spectrum in the roots and capsules of the species and hybrids of *Papaver* section *Oxytona* 563
Miura, H., see Goto, J. 119
Montzka, S. A., see McNamara, E. A. 75
Mori, S.
Interchangeable use of aqueous and organic solvents in a hydrophilic poly(vinyl alcohol) gel column 137
Mutti, A., see Ghiggeri, G. M. 347
Myher, J. J.
——, Kuksis, A. and Marai, L.
Identification of the more complex triacylglycerols in bovine milk fat by gas chromatography–mass spectrometry using polar capillary columns 93
Nagakura, R., see Goto, J. 119
Nagatsu, R., see Niwa, T. 85
Nakayama, F.
Quantitative microanalysis of bile acids in biological samples. Collaborative study 399
Nambara, T., see Goto, J. 119

Nelis, H. J. C. F.
—— and De Leenheer, A. P.
Reversed-phase liquid chromatography of astacene 535
Nes, W. D., see Xu, S. 377
Niwa, T.
——, Takeda, N., Tatematsu, A., Matsuura, S., Yoshida, M. and Nagatsu, R.
Migration of tetrahydroisoquinoline, a possible parkinsonian neurotoxin, into monkey brain from blood as proved by gas chromatography–mass spectrometry 85
Norton, R. A., see Xu, S. 377
Nováková, N., see Janča, J. 549
Novotny, M., see David, P. A. 623
Nyéki, O., see Görög, S. 317
O'Flynn, M. A.
——, Causon, R. C., Brown, J. and Kageyama, S.
Development of a solid-phase extraction technique for α-human atrial natriuretic peptide in human plasma 469
Palevitsch, D., see Milo, J. 563
Palma, P., see Mangani, F. 527
Patel, P., see Jack, D. M. 257
Peichang, L.
—— and Hongxin, H.
Expert system for chromatography 175
Ponomareva, T. M., see Davankov, V. A. 309
Poole, C. F., see Kersten, B. R. 191
Ranný, M., see Michalec, Č. 543
Rassi, Z. El, see Maa, Y.-F. 331
Righetti, P.G., see Ghiggeri, G.M. 347
Rooney, L., see Jack, D. B. 257
Saavedra, J. E., see Issaq, H. J. 511
Sagliano, Jr. N., see Floyd, T. R. 43
Satterwhite, D. M.
—— and Croteau, R. B.
Applications of gas chromatography to the study of terpenoid metabolism 61
Schaeffler, I., see Weissenberg, M. 485
Schilt, R., see Farjam, A. 419
Schön, I., see Görög, S. 317
Schultz, N., see Issaq, H. J. 511
Schwedt, G., see Golombek, F. 283
Shohat, S.
——, Grushka, E. and Glikberg, S.
Liquid chromatographic analysis of organic additives in copper plating baths 503
Sievers, R. M., and see McNamara, E. A. 75
Sjövall, J., see Marshall, H.-U. 459
Smith, R. J., see Haken, J. K. 31
Smolková-Keulemansová, E., see Snopek, J. 571
Smrz, M., see Hanis, T. 443
Snopek, J.
——, Jelínek, I. and Smolková-Keulemansová. E.
Micellar, inclusion and metal-complex enantioselective pseudophases in high-performance electromigration methods 571
Stafford, A. E., see Lin, J.-T. 519
Suzuki, H., see Goto, J. 119
Takeda, N., see Niwa, T. 85
Tatematsu, A., see Niwa, T. 85
Tchapla, A., see Martin, M. 157
Thevenon, G., see Martin, M. 157
Tipker, J.
——, Groen, C. P., Van den Bergh-Swart, J. K. and Van den Berg, J. H. M.
Contribution of electronic effects to the lipophilicity determined by comparison of values of log *P* obtained by high-performance liquid chromatography and calculation 227
Trainor, T. M.
——, Giese, R. W. and Vouros, P.
Mass spectrometry of electrophore-labeled nucleosides pentafluorobenzyl and cinnamoyl derivatives 369
Turner, R., see Jackson, P. J. 359
Van den Berg, J. H. M., see Tipker, J. 227
Van den Bergh-Swart, J. K., see Tipker, J. 227
Verheggen, Th. P. E. M.
——, Beckers, J. L. and Everaerts, F. M.
Simple sampling device for capillary isotachophoresis and capillary zone electrophoresis 615
——, see Beckers, J. L. 591
Vouros, P., see Trainor, T. M. 369
Wall, F., see Huang, T. 409
Warner, F. P., see Dawkins, J. V. 145
Weissenberg, M.
——, Levy, A., Schaeffler, I. and Levy, E. C.
High-performance liquid chromatographic analysis of the ajmalicine distribution in roots of *Catharanthus roseus* lines with different flower colours 485
Westerlund, D., see Johansson, M. 241
Williams, D. G., see Issaq, H. J. 511
Xu, S.
——, Norton, R. A., Crumley, F. G. and Nes, W. D.
Comparison of the chromatographic properities of sterols, select additional steroids and triterpenoids: gravity-flow column liquid chromatography, thin-layer chromatography, gas–liquid chromatography and high-performance liquid chromatography 377
Yoshida, M., see Niwa, T. 85